

University of Groningen

Staphylococcus aureus spa type t437

Glasner, C; Pluister, G; Westh, H; Arends, J P; Empel, J; Giles, E; Laurent, F; Layer, F; Marstein, L; Matussek, A

Published in:
Clinical Microbiology and Infection

DOI:
[10.1016/j.cmi.2014.09.010](https://doi.org/10.1016/j.cmi.2014.09.010)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Glasner, C., Pluister, G., Westh, H., Arends, J. P., Empel, J., Giles, E., Laurent, F., Layer, F., Marstein, L., Matussek, A., Mellmann, A., Pérez-Vásquez, M., Ungvári, E., Yan, X., Zemličková, H., Grundmann, H., & van Dijk, J. M. (2015). Staphylococcus aureus spa type t437: identification of the most dominant community-associated clone from Asia across Europe. *Clinical Microbiology and Infection*, 21(2), 163.e1-163.e8. <https://doi.org/10.1016/j.cmi.2014.09.010>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Staphylococcus aureus spa type t437: identification of the most dominant community-associated clone from Asia across Europe

C. Glasner¹, G. Pluister², H. Westh^{3,4}, J. P. Arends¹, J. Empel⁵, E. Giles⁶, F. Laurent⁷, F. Layer⁸, L. Marstein⁹, A. Matussek¹⁰, A. Mellmann¹¹, M. Pérez-Vásquez¹², E. Ungvári¹³, X. Yan^{14,15}, H. Žemličková¹⁶, H. Grundmann¹ and J. M. van Dijl^{1*}

1) Department of Medical Microbiology, University of Groningen and University Medical Center Groningen, Groningen, The Netherlands, 2) Bacterial Surveillance and Response, Center for Infectious Disease Control, National Institute of Public Health and the Environment, Bilthoven, The Netherlands, 3) Department of Clinical Microbiology, Hvidovre Hospital, Hvidovre, Denmark, 4) Faculty of Medical Sciences, University of Copenhagen, Copenhagen, Denmark, 5) Department of Molecular Microbiology National Medicines Institute, Warsaw, Poland, 6) Department of Microbiology, Scottish MRSA Reference Laboratory, Glasgow, United Kingdom, 7) Centre National de Référence des Staphylocoques, Université de Lyon, INSERM U851, Lyon, France, 8) National Reference Centre for Staphylococci and Enterococci, Division Nosocomial Pathogens and Antibiotic Resistances, Department of Infectious Diseases, Robert Koch Institute, Wernigerode, Germany, 9) Department of Medical Microbiology, MRSA Reference Laboratory, St. Olavs Hospital, Trondheim University Hospital, Norway, 10) Department of Laboratory Services, County Hospital Ryhov, Jönköping, Sweden, 11) Institute for Hygiene University Hospital Münster, Münster, Germany, 12) Servicio de Bacteriología, Centro Nacional de Microbiología, Instituto de Salud Carlos III, Majadahonda, Madrid, Spain, 13) Department of Phage Typing and Molecular Epidemiology, National Center for Epidemiology, Budapest, Hungary, 14) State Key Laboratory for Infectious Disease Prevention and Control, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China, 15) Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, Hangzhou, China and 16) National Institute of Public Health, Prague, Czech Republic

Abstract

Methicillin-resistant *Staphylococcus aureus* (MRSA) belonging to the multilocus sequence type clonal complex 59 (MLST CC59) is the predominant community-associated MRSA clone in Asia. This clone, which is primarily linked with the *spa* type t437, has so far only been reported in low numbers among large epidemiological studies in Europe. Nevertheless, the overall numbers identified in some Northern European reference laboratories have increased during the past decade. To determine whether the *S. aureus* t437 clone is present in other European countries, and to assess its genetic diversity across Europe, we analysed 147 *S. aureus* t437 isolates from 11 European countries collected over a period of 11 years using multiple locus variable number tandem repeat fingerprinting/analysis (MLVF/MLVA) and MLST. Additionally 16 *S. aureus* t437 isolates from healthy carriers and patients from China were included. Most isolates were shown to be monophyletic with 98% of the isolates belonging to the single MLVA complex 621, to which nearly all included isolates from China also belonged. More importantly, all MLST-typed isolates belonged to CC59. Our study implies that the European *S. aureus* t437 population represents a genetically tight cluster, irrespective of the year, country and site of isolation. This underpins the view that *S. aureus* CC59 has been introduced into several European countries, not being restricted to particular geographical regions or specific host environments. The European *S. aureus* t437 isolates thus bear the general hallmarks of a high-risk clone.

Clinical Microbiology and Infection © 2014 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

Keywords: Asia, CC59, community-associated, Europe, identification, MRSA, *spa* type t437, *Staphylococcus aureus*

Original Submission: 25 June 2014; **Revised Submission:** 12 August 2014; **Accepted:** 11 September 2014

Editor: G. Lina

Article published online: 29 October 2014

Corresponding author: J.M. van Dijl, Department of Medical Microbiology, University of Groningen, University Medical Center Groningen, Hanzeplein 1, P.O. Box 30001, 9700 RB Groningen, The Netherlands

E-mail: j.m.van.dijl01@umcg.nl
These authors contributed equally to this work

Introduction

The emergence and spread of human pathogens, such as *Staphylococcus aureus*, among hospital patients as well as in the community are threatening public health worldwide. The capability to acquire antibiotic resistance and a plethora of virulence factors make *S. aureus* formidably apt to cause disease in these different settings. This is underscored by the large number of different *S. aureus* types encountered in many hosts and environments [1–6].

Epidemiological and, more recently, molecular studies have shown that certain clones of *S. aureus* attain a geo-spatial predominance [1–6]. Importantly, various community-associated methicillin-resistant *S. aureus* (CA-MRSA) clones have evolved independently on different continents. Multilocus sequence type (MLST/ST) 80 is the predominant CA-MRSA clone in Europe, ST93 in Australia, ST30 in Oceania, ST8 in the United States of America (USA), and ST59 in Asia. Nevertheless, the exchange of clones between countries and continents has been observed [3,4,7,8,9], as can be expected from the current reach, volume and speed of travel [10].

The ST59 clone, which is the founder of the MLST clonal complex 59 (CC59), is one of the most frequent multidrug resistant CA-MRSA clones in Asia [8]. In 2007, Tristan and colleagues reported for the first time Asian *S. aureus* ST59 isolates in association with the *spa* type t437 [11]. Subsequently, a large community and hospital study across Asia described the CC59 as the most prevalent CC, including ST59 and its variants ST1241 and ST338; moreover, ST59-MRSA-t437 was identified as the most prevalent clone between 2004 and 2006 [8]. In the study by Song *et al.*, the collected CA- and hospital-associated MRSA CC59 isolates from Asia showed rapid spread between hospitals and the community in a bi-directional manner, and also across borders [8]. A similar picture emerged from other studies in Asia and Western Australia where ST59-MRSA-t437 was identified as a major clone [12–16]. In contrast, *S. aureus* CC59 isolates were incidentally reported in the USA [17] and in Europe [18–22].

Recently, a study from Belgium identified four (1%) ST338-t437 CA-MRSA isolates amongst 410 MRSA isolates collected between 2005 and 2009 [23]. Furthermore, a multicentre study performed in the 16 most populous European countries identified a total number of 22 (6%) *S. aureus* CC59 isolates with the *spa* type t437 of which one (4.5%) was methicillin-sensitive *S. aureus* (MSSA) [24]. Intriguingly, Rolo and co-workers reported an increased frequency of the ST59 clone since 2007, and they concluded that this clone was most prevalent in Northern Europe (Finland, Sweden and Poland) [24]. This is in line with our own observations that the numbers of *S. aureus* t437 isolates

in Norway, Denmark and Poland are increasing. Specifically, in Norway *S. aureus* t437 is among the most commonly identified MRSA clones corresponding to ~2.5% of all MRSA isolates every year since 2008 (L. Marstein, personal communication). In Denmark, the numbers of *S. aureus* t437 isolates have gone up from zero before 2006, through 1–3 cases per year between 2006 and 2008, to 7–14 cases per year since 2007 (H. Westh, personal communication). Lastly, the ST338-t437 PVL-positive clone seems currently to be the most prevalent CA-MRSA clone in Poland (J. Empel, personal communication).

Altogether, the combined literature data for Asia and Europe, and the apparently increasing numbers of *S. aureus* t437 isolates in Northern Europe formed the incentive to assess the presence of *S. aureus* t437 in other European countries and, more importantly, to determine their genetic relatedness. Clearly, the perceived risk for dissemination and establishment of a new community-associated *S. aureus* clone with a multidrug resistant phenotype in Europe would justify appropriate preventive infection control measures.

Materials and methods

Bacterial isolates

A total of 163 *S. aureus* isolates (147 from Europe and 16 from China) with the *spa* type t437, collected between 2002 and 2012 from patients and healthy carriers in 11 different European countries and four different geographical locations in China (i.e. Hangzhou, Zhejiang Province; Hefei, Anhui Province; Harbin, Heilongjiang Province; and Beijing), were analysed in the present study. The epidemiological and molecular characteristics of the isolates, including origin, year of isolation, antibiotic phenotype and information on the patients are presented in Appendix Table A1 in the supporting information. One isolate from the Czech Republic with an unrelated *spa* type t442 was also included in the present study.

Antibiotic susceptibility and presence of *mecA* and the PVL locus

Antibiotic susceptibility for benzylpenicillin, chloramphenicol, ciprofloxacin, clindamycin (constitutive), erythromycin, fosfomycin, fusidic acid, gentamicin, kanamycin, linezolid, mupirocin, oxacillin, rifampicin, teicoplanin, tetracycline, tobramycin, trimethoprim/sulfamethoxazole and vancomycin was determined with the VITEK 2 system (AST P633 card, bioMérieux, Marcy l'Etoile, France). The VITEK 2 minimum inhibitory concentration results were interpreted using the Advanced Expert System following EUCAST guidelines (www.eucast.org). The presence of the *mecA* and PVL-encoding genes (*lukF-PV* and *lukS-PV*) was determined by PCR [25].

Extraction of total DNA for typing

Total DNA for the multiple-locus variable number tandem repeat fingerprinting (MLVF) typing was prepared as previously described [2]. The preparation of lysates for multiple locus variable number tandem repeat analysis (MLVA) and *spa* typing was performed as described by Schouls et al. [25].

spa typing

The *spa* typing was performed according to the protocol previously described [26]. The *spa* types were assigned through the use of Ridom StaphType software version 2.2.1 (Ridom GmbH, Münster, Germany) and the SpaServer (<http://www.spaserver.ridom.de>).

Multiple-locus variable number tandem repeat fingerprinting

MLVF was performed as described by Glasner et al. and Sabat et al. using the Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA) to separate PCR fragments [2,27]. For the analysis of electropherograms with Gel Compar II (Applied Maths, Kortrijk, Belgium), the position tolerance and optimization were set to 0.5% and 0.5%, respectively, and the dice formula was used to calculate the pairwise similarity coefficient. With the selected position tolerance, all Bioanalyzer runs for the control isolate M2 (clustering together at the bottom part of the MLVF dendrogram) were identical [27]. A dendrogram was created with the unweighted pair-group method using geometric averages (UPGMA).

Multiple-locus variable number tandem repeat analysis

MLVA was performed according to Schouls et al. [25]. Isolates that differed by one or more alleles were considered distinct types. Minimum spanning tree analysis of MLVA was performed using BioNumerics software (Applied Maths, Kortrijk, Belgium) to group related MLVA types (MTs) into MLVA complexes (MCs). Such MCs encompass single locus variants as described by Schouls et al. [25]. A singleton was defined as an MT that was not grouped into an MC.

Multilocus sequence typing

MLST was performed on representative *S. aureus* t437 isolates of each MT as described by Enright et al. [28]. In brief, the allelic profiles of each selected isolate were obtained by sequencing internal fragments of seven housekeeping genes (*arcC*, *aroE*, *glpF*, *gmk*, *pta*, *tpi*, *yqiL*) and entering them on the MLST homepage (<http://saureus.mlst.net>), where seven numbers depicting the allelic profile were assigned that defined a ST [28]. The allelic profiles of the *S. aureus* t437 isolates were compared by using the based upon related sequence types (BURST) program [4].

Results

Collection of *S. aureus* t437 isolates from Europe and China

To explore the presence and genetic relatedness of *S. aureus* t437 across Europe, a convenience sample of 147 isolates with the *spa* type t437 was established at the Medical Microbiology Department at the University Medical Center Groningen (UMCG, The Netherlands). This was achieved through the identification of *S. aureus* *spa* type t437 isolates by inspection of the RIDOM *spa* server database (www.spaserver.ridom.de). In addition, representatives from all staphylococcal expert and reference laboratories were approached and asked for *S. aureus* *spa* type t437 isolates. All *S. aureus* isolates were then sent to the UMCG, stored and subsequently propagated for subsequent molecular analyses. The *spa* type t437 is currently ranked 24th on the RIDOM *spa* server with a frequency of 0.63% and a total number of 1878 isolates (June 2014). The collected *S. aureus* t437 isolates were from 11 European countries, namely the Netherlands ($n = 64$), Scotland ($n = 27$), Norway ($n = 20$), Germany ($n = 13$), Denmark ($n = 10$), Spain ($n = 3$), Poland ($n = 3$), France ($n = 3$), Sweden ($n = 2$), Czech Republic ($n = 1$) and Hungary ($n = 1$). Sixteen *S. aureus* isolates with the *spa* type t437 from patients and healthy carriers from China that were available during the time of investigation at our institute were added to the collection. One isolate from the Czech Republic was originally submitted as *S. aureus* *spa* type t437 but, during the course of investigation, it was determined to have the unrelated *spa* type t442. Seventy-eight (47.9%) *S. aureus* t437 isolates were sampled from abscesses and skin infections, and 56 (34.3%) were isolated from blood or the nose, ear or throat (supporting information Appendix Table A1). The source of isolation is unknown for the remaining 29 (17.8%) *S. aureus* t437 isolates.

Antibiotic resistance and proportion of *pvl* and *mecA*

Antibiotic susceptibility testing showed that the antibiotic profiles of the 143 (87.7%) MRSA t437 isolates were very similar to those of the 20 (12.3%) MSSA t437 isolates (Table I). Consistent distinctions in the resistance of the investigated MRSA t437 and MSSA t437 isolates were only observed for oxacillin and chloramphenicol. The increased chloramphenicol resistance of the collected MRSA t437 isolates is in accordance with results from Asian CA-MRSA studies [7,15]. The majority of the investigated t437 isolates were resistant to clindamycin (constitutive), erythromycin, kanamycin and penicillin, and all t437 isolates were susceptible to fosfomycin, fusidic acid, linezolid, mupirocin, rifampicin, teicoplanin and vancomycin (Table I). No clear association between countries of origin or

TABLE 1. Antibiotic resistance profiles of the 143 MRSA t437 and 20 MSSA t437 isolates*

Antibiotic	MRSA	MSSA
	n (resistance rate, %)	n (resistance rate, %)
Ciprofloxacin	2 (1.4)	2 (10)
Chloramphenicol	93 (65)	3 (15)
Clindamycin (constitutive)	118 (82.5)	17 (85)
Erythromycin	121 (84.6)	17 (85)
Fosfomycin	0 (0)	0 (0)
Fusidic acid	0 (0)	0 (0)
Gentamicin	3 (2.1)	2 (10)
Kanamycin	120 (83.9)	16 (80)
Linezolid	0 (0)	0 (0)
Mupirocin	0 (0)	0 (0)
Oxacillin	143 (100)	0 (0)
Penicillin	143 (100)	18 (90)
Rifampicin	0 (0)	0 (0)
Teicoplanin	0 (0)	0 (0)
Tetracycline	102 (71.3)	11 (55)
Tobramycin	3 (2.1)	2 (10)
Trimethoprim/sulfamethoxazole	1 (0.7)	5 (25)
Vancomycin	0 (0)	0 (0)

MRSA, methicillin-resistant *Staphylococcus aureus*; MSSA, methicillin-sensitive *S. aureus*.

*For details on the antibiotic resistances of the study isolates, please see supporting information [Appendix Table A1](#).

sample source and a certain antibiotic resistance profile could be detected. In accordance with the observed oxacillin resistance profiles, 88% of the isolates tested positive for *mecA* (143 isolates). Furthermore, 82% (134 isolates) tested positive for the Panton-Valentine leukocidin (PVL) locus [3]. The majority of *S. aureus* t437 isolates were thus both *mecA*- and *pvl*-positive (127 isolates, 78%), as previously reported for Asian CC59 isolates [12,15]. Fifteen isolates were *mecA*-positive and *pvl*-negative, while 21 isolates were *mecA*-negative of which 8 were *pvl*-positive and 13 *pvl*-negative.

MLVF

MLVF analysis identified 37 different banding patterns among the 163 *S. aureus* t437 isolates as shown in [Fig. 1](#). Sixteen patterns were represented by two or more isolates (142 isolates in total). The remaining 21 patterns each consisted of a single isolate. Application of previously published cut-off values of 64%, 67% or 75% led to 2, 2 and 9 clusters, respectively [2,27]. The two lower cut-off values joined 161 and 154 isolates into one major cluster respectively, indicating a high genomic relatedness of these isolates. Even without the application of a cut-off value resulting in MLVF clusters, the relatedness of all *S. aureus* t437 isolates can be inferred by inspection from the highly similar MLVF banding patterns (see also [2,27]). Slight differences in the band sizes indicate loss or gain of repeats in the respective variable number of tandem repeats (VNTRs). As expected, the isolate with the unrelated *spa* type t442 displayed a different MLVF pattern with only 34% similarity to the closest related isolate in the MLVF dendrogram ([Fig. 1](#)). The MLVF patterns did not unveil any epidemiological signal, such as the country of

origin, year of isolation or source. Thus isolates from different countries, even from China, appeared randomly distributed over the MLVF dendrogram. Notably, MLVF is a highly discriminatory but non-portable PCR-based DNA fingerprinting method that utilizes size differences in five coding regions (*spa*, *sspA*, *sdnCDE*, *clfA* and *clfB*) containing VNTRs. In contrast, the subsequently implemented MLVA method for DNA typing is suitable for inter-laboratory comparisons and allows the determination of clonal relationships between isolates. To this end, MLVA targets ten non-coding loci, which are sequenced and for which the exact numbers of repeat units are measured. This approach is therefore more precise than the visualization of MLVF results on agarose gels or microfluidic chips.

MLVA

The 163 *S. aureus* t437 isolates produced 13 different MTs, namely MT621, MT1035, MT1297, MT1831, MT1870, MT1875, MT2075, MT2322, MT3560, MT4124, MT4125, MT4126 and MT4183 ([Fig. 2A](#) and supporting information [Appendix Table A1](#)). The MT621 was clearly the most predominant type in the present collection comprising 133 (82%) isolates and, more importantly, 159 (98%) *S. aureus* isolates belonged to the same MC0621, comprising 9 different MTs ([Fig. 2A](#)). The remaining four isolates were MLVA singletons (MCnone). Generally, six MTs were shared by 2 or more isolates (156 isolates in total), whereas 7 MTs were represented by single isolates. The included t422 isolate from the Czech Republic belonged to the unrelated MT165 and MC5.

MLST

A total of 46 representative *S. aureus* t437 isolates comprising one MT each were selected for MLST analysis ([Fig. 2B](#)). This showed that 39 (85%) of these isolates belonged to the CC59. ST59 was the predominant ST with 26 (57%) isolates, and the two single-locus variants ST87 and ST338 were identified 2 (4%) and 11 (24%) times, respectively. The remaining 7 isolates, which were all from China, belonged to ST2147 differing from ST59 by three alleles. The included isolate with the *spa*-type t422 belonged to the unrelated ST5. The relationship between the MLST and MLVA data is depicted in [Fig. 2B](#). The MLST minimum spanning tree shows that ST59, the founder of CC59, is composed of 10 different MTs.

Discussion

The present study underpins the view that the *S. aureus* ST59 clone and other STs of CC59 with the *spa* type t437 which are commonly encountered in Asia, are present in several European countries. More importantly, by implementing three different




FIG. 1. MLVF dendrogram of the 163 investigated *S. aureus* t437 isolates generated by the UPGMA algorithm. In addition to the study isolates, one isolate with a different *spa* type (t442), MLVA type and MLST type, and 16 controls (designated M2) were included in the analysis. Details on the different isolates from top to bottom are provided in the same order in [Appendix Table A1](#) in the supporting information. MLVA, multiple-locus variable number tandem repeat analysis; MLST, multilocus sequence typing; MLVF, multiple locus variable number tandem repeat fingerprinting; UPGMA, unweighted pair-group method using geometric averages. *Lanes corresponding to Chinese t437 isolates.




FIG. 2. (A) Minimum spanning tree of the 163 *S. aureus* t437 isolates typed by MLVA. Clustering of MLVA profiles was performed with a categorical coefficient. In the minimum spanning tree, the MLVA types are displayed as circles labelled with different colours. The size of each circle indicates the number of isolates of the particular MLVA type. MLVA complexes were assigned if two neighbouring types did not differ in more than one VNTR locus. MLVA types and complexes are indicated in characters, e.g. 621 denotes MLVA type 621, and MC0621 denotes MLVA complex 621. The three MTs 1870, 4125 and 4126 are double-locus variants of MT0621 and do not belong to any MC (MCnone). The t442 isolate from the Czech Republic is also included in the minimum spanning tree, showing that it belongs to the unrelated MT165. (B) MLST analysis of 46 representative *S. aureus* t437 isolates with BURST identifies one MLST CC59, which includes ST59 (26), ST338 (10) and ST87 (2). The two other identified STs, ST2147 (7) and ST5 (one isolate from the Czech Republic with *spa* type t442) are singletons. The colours within each ST circle indicate the different MTs belonging to the respective ST. MLST, multilocus sequence typing; MLVA, multiple-locus variable number tandem repeat analysis; MT, MLVA types; ST, sequence typing; VNTR, variable number of tandem repeats.

highly discriminatory molecular typing tools, namely MLVF, MLVA and MLST, we demonstrate a high degree of molecular similarity in the studied *S. aureus* t437 isolates that were collected over a period of 10 years from 11 different European countries. This shows that this specific *S. aureus* clone is not restricted to particular geographical regions or specific host environments, and that *S. aureus* t437 in Europe belongs to a very tight molecular cluster of *S. aureus* isolates. The European *S. aureus* t437 isolates thus bear the general hallmarks of a high-risk clone.

At least 35 (21.5%) of the patients from whom the currently investigated European *S. aureus* t437 isolates had been collected were immigrants or adopted children, or had travelled to countries outside Europe, Asia in particular (supporting

information [Appendix Table A1](#); note that for 104 isolates no such information could be retrieved). Such patients may have introduced the CC59 clone in Europe and subsequently transmitted it to other European citizens, including their family members, as can be inferred from the analysis of the isolates from the Netherlands and Denmark (supporting information [Appendix Table A1](#)). This view is also supported by the high genetic relatedness of the European isolates with the 16 Chinese *S. aureus* t437 isolates included in our study as shown by MLVF and MLVA. Although seven Chinese isolates belonged to ST2147 (a triple-locus variant of ST59) and only one isolate belonged to ST59, the Chinese isolates clustered among all other *S. aureus* t437 isolates in the MLVF dendrogram ([Fig. 1](#) and supporting

information Appendix Table A1), and 15 of these isolates belonged to the MC0621 which includes 98% of the 163 investigated *S. aureus* t437 isolates. Notably, in most Asian studies only MRSA isolates were collected and typed. It is therefore conceivable that MSSA with the *spa*-type t437 belonging to CC59 has thus far been overlooked in Asia. In contrast, the present European *S. aureus* t437 isolates also include MSSA isolates. This observation is in line with the findings reported by Rolo et al. [24]. Otherwise, the antibiotic resistance profiles of the presently investigated European *S. aureus* t437 isolates were very similar to those described for Asian CC59 isolates, but not to those of other CA *S. aureus* clones from Europe [3,8,14,29,30]. Lastly, the frequency of the PVL-encoding genes (82%) in the present *S. aureus* t437 sample is in accordance with the reported numbers in the Asian studies [12,15,31].

In conclusion, we have combined MLVF, MLVA and MLST to obtain a high-resolution snapshot of the *S. aureus* t437 population in Europe based on retrospectively collected isolates that had already been *spa* typed. Since *spa* typing or any of the three other molecular typing methods implemented in the present study are not yet performed on a daily routine basis in most local laboratories and hospitals, we consider it likely that the CC59 clone has so far remained under-detected in Europe. This is a cause for concern in view of the predominance of *S. aureus* CC59 in Asia and its clinical repercussions, in particular multi-drug resistant soft tissue and skin infections, even though the prevalence of CC59 in Europe is probably still relatively low. Where possible, the further dissemination of this potentially high-risk clone should therefore be prevented, for example through active screening of patients with staphylococcal infections who have a history of travel in Asia.

Transparency declaration

Funding: C.G. was supported by a fellowship from the Graduate School for Medical Sciences of the University of Groningen. J.E. was partly supported by the Ministry of Health within the framework of the Module I of the National Programme of Antibiotic Protection and by the Ministry of Science and Higher Education (Mikrobank 2 Programme, and a grant no 1216/7. PR UE/2009/7).

Conflict of interest: The authors declare no conflicts of interest.

Author contributions

C. Glasner, H. Grundmann and J. M. van Dijk: designed the study and wrote the manuscript. C. Glasner and G. Pluister:

performed laboratory investigations. G. Pluister, H. T. Westh, J. P. Arends, J. Empel, E. Giles, F. Laurent, F. Layer, L. Marstein, A. Matussek, A. Mellmann, M. Perez-Vasquez, E. Ungvári, X. Yan and H. Zemlickova: performed epidemiological investigations, provided the study isolates, provided feedback, contributed with comments and reviewed the manuscript.

Acknowledgements

The authors thank the staff of the Bacteriology Laboratory at the UMCG for their support in determining the antibiotic resistance profiles. We would also like to thank Dr. Kit Boye, Bonnie Cosgrove, Aleksandra Kozińska, Helene Meugnier, Dr. Akós Tóth, Dr. Ana Vindel and Professor Dr Jianzhong Zhang for the provision of clinical isolates.

Appendix A. Supplementary data

Supplementary data related to this article can be found online at <http://dx.doi.org/10.1016/j.cmi.2014.09.010>.

References

- [1] Glasner C, Sabat AJ, Chlebowicz MA, Vanderhaeghen W, Fetsch A, Guerra B, et al. High-resolution typing by MLVF unveils extensive heterogeneity of European livestock-associated methicillin-resistant *Staphylococcus aureus* isolates with the sequence type 398. *Int J Med Microbiol* 2013;303:124–7.
- [2] Glasner C, Sabat AJ, Dreisbach A, Larsen AR, Friedrich AW, Skov RL, et al. Rapid and high-resolution distinction of community-acquired and nosocomial *Staphylococcus aureus* isolates with identical pulsed-field gel electrophoresis patterns and *spa* types. *Int J Med Microbiol* 2013;303:70–5.
- [3] Vandenesch F, Naimi T, Enright MC, Lina G, Nimmo GR, Heffernan H, et al. Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Panton-Valentine leukocidin genes: worldwide emergence. *Emerg Infect Dis* 2003;9:978–84.
- [4] Enright MC, Robinson DA, Randle G, Feil EJ, Grundmann H, Spratt BG. The evolutionary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proc Natl Acad Sci USA* 2002;99:7687–92.
- [5] Holden MT, Hsu LY, Kurt K, Weinert LA, Mather AE, Harris SR. A genomic portrait of the emergence, evolution, and global spread of a methicillin-resistant *Staphylococcus aureus* pandemic. *Genome Res* 2013;23:653–64.
- [6] Grundmann H, Aanensen DM, van den Wijngaard CC, Spratt BG, Harmsen D, Friedrich AW, et al. Geographic distribution of *Staphylococcus aureus* causing invasive infections in Europe: a molecular-epidemiological analysis. *PLoS Med* 2010;7:e1000215.
- [7] Takano T, Higuchi W, Zaraket H, Otsuka T, Baranovich T, Enany S, et al. Novel characteristics of community-acquired methicillin-resistant *Staphylococcus aureus* strains belonging to Multilocus Sequence Type 59 in Taiwan. *Antimicrob Agents Chemother* 2008;52:837–45.
- [8] Song JH, Hsueh PR, Chung DR, Ko KS, Kang CI, Peck KR, et al. Spread of methicillin-resistant *Staphylococcus aureus* between the community

- and the hospitals in Asian countries: an ANSORP study. *J Antimicrob Chemother* 2011;66:1061–9.
- [9] DeLeo FR, Otto M, Kreiswirth BN. Community-associated methicillin-resistant *Staphylococcus aureus*. *Lancet* 2010;375:1557–68.
- [10] Tatem AJ, Rogers DJ, Hay SI. Global transport networks and infectious disease spread. *Adv Parasitol* 2006;62:293–343.
- [11] Tristan A, Bes M, Meugnier H, Lina G, Bozdogan B, Courvalin P, et al. Global distribution of Pantone-Valentine Leukocidin-positive methicillin-resistant *Staphylococcus aureus*, 2006. *Emerging Infect Dis* 2007;13:594–7.
- [12] Ho CM, Ho MW, Lee CY, Tien N, Lu JJ. Clonal spreading of methicillin-resistant SCCmec *Staphylococcus aureus* with specific *spa* and *dru* types in central Taiwan. *Eur J Clin Microbiol Infect Dis* 2012;31:499–504.
- [13] Coombs GW, Monecke S, Ehrlich R, Slickers P, Pearson JC, Tan HL, et al. Differentiation of clonal complex 59 community-associated methicillin-resistant *Staphylococcus aureus* in Western Australia. *Antimicrob Agents Chemother* 2010;54:1914–21.
- [14] Ko KS, Lee JY, Suh JY, Oh WS, Peck KR, Lee NY, et al. Distribution of major genotypes among methicillin-resistant *Staphylococcus aureus* clones in Asian countries. *J Clin Microbiol* 2005;43:421–6.
- [15] Wu D, Wang Q, Yang Y, Geng W, Wang Q, Yu S, et al. Epidemiology and molecular characteristics of community-associated methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* from skin/soft tissue infections in a children's hospital in Beijing, China. *Diagn Microbiol Infect Dis* 2010;67:1–8.
- [16] Ho PL, Chuang SK, Choi YF, Lee RA, Lit AC, Ng TK, et al. Community-associated methicillin-resistant and methicillin-sensitive *Staphylococcus aureus*: skin and soft tissue infections in Hong Kong. *Diagn Microbiol Infect Dis* 2008;61:245–50.
- [17] Pan ES, Diep BA, Charlebois ED, Auerswald C, Carleton HA, Sensabaugh GF, et al. Population dynamics of nasal strains of methicillin-resistant *Staphylococcus aureus* - and their relation to community-associated disease activity. *J Infect Dis* 2008;192:811–8.
- [18] Hedin G, Fang H. Epidemiology of methicillin-resistant *Staphylococcus aureus* in Southern Stockholm, 2000–2003. *Microb Drug Resist* 2007;13:241–50.
- [19] Monecke S, Jatzwauk L, Weber S, Slickers P, Ehrlich R. DNA microarray-based genotyping of methicillin-resistant *Staphylococcus aureus* strains from Eastern Saxony. *Clin Microbiol Infect* 2008;14:534–45.
- [20] Huijsdens XW, van Santen-Verheul MG, Spalburg E, Heck ME, Pluister GN, Eijkelkamp BA, et al. Multiple cases of familial transmission of community-acquired methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 2006;44:2994–6.
- [21] Łuczak-Kadłubowska A, Sulikowska A, Empel J, Piasecka A, Orczykowska M, Kozinska A, et al. Countrywide molecular survey of methicillin-resistant *Staphylococcus aureus* strains in Poland. *J Clin Microbiol* 2008;46:2930–7.
- [22] Ellington MJ, Perry C, Ganner M, Warner M, McCormick Smith I, Hill RL, et al. Clinical and molecular epidemiology of ciprofloxacin-susceptible MRSA encoding PVL in England and Wales. *Eur J Clin Microbiol Infect Dis* 2009;28:1113–21.
- [23] Brauner J, Hallin M, Deplano A, De Mendonça R, Nonhoff C, De Ryck R, et al. Community-acquired methicillin-resistant *Staphylococcus aureus* clones circulating in Belgium from 2005 to 2009: changing epidemiology. *Eur J Clin Microbiol Infect Dis* 2012;32:613–20.
- [24] Rolo J, Miragaia M, Turlej-Rogacka A, Empel J, Bouchami O, Faria NA, et al. High genetic diversity among community-associated *Staphylococcus aureus* in Europe: results from a multicenter study. *PLoS ONE* 2012;7:e34768.
- [25] Schouls LM, Spalburg EC, van Luit M, Huijsdens XW, Pluister GN, van Santen-Verheul MG, et al. Multiple-locus variable number tandem repeat analysis of *Staphylococcus aureus*: comparison with pulsed-field gel electrophoresis and *spa*-typing. *PLoS ONE* 2009;4:e5082.
- [26] Mellmann A, Friedrich AW, Rosenkötter N, Rothgänger J, Karch H, Reintjes R, et al. Automated DNA sequence-based early warning system for the detection of methicillin-resistant *Staphylococcus aureus* outbreaks. *PLoS Med* 2006;3:e33.
- [27] Sabat AJ, Chlebowicz MA, Grundmann H, Arends JP, Kampinga G, Meessen NE, et al. Microfluidic chip-based multiple-locus variable-number tandem repeat fingerprinting (MLVF) with new primer sets for methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 2012;50:2255–62.
- [28] Enright MC, Day NPJ, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 2000;38:1008–15.
- [29] Huang YC, Hwang KP, Chen PY, Chen CJ, Lin TY. Prevalence of methicillin-resistant *Staphylococcus aureus* nasal colonization among Taiwanese children in 2005 and 2006. *J Clin Microbiol* 2007;45:3992–5.
- [30] Wang C-C, Lo W-T, Chu M-L, Siu LK. Epidemiological typing of community-acquired methicillin-resistant *Staphylococcus aureus* isolates from children in Taiwan. *Clin Infect Dis* 2004;39:481–7.
- [31] Chen CJ, Su LH, Chiu CH, Lin TY, Wong KS, Chen YY, et al. Clinical features and molecular characteristics of invasive community-acquired methicillin-resistant *Staphylococcus aureus* infections in Taiwanese children. *Diagn Microbiol Infect Dis* 2007;59:287–93.